

Dr. Pamela Hawranik RN PhD, Associate Professor, Dean, Faculty of Graduate Studies, Athabasca University
Sandy Bell RN MN, Director Quality, Patient Safety & Education Services, Misericordia Health Centre

Acknowledgements: Funding was awarded by Fort Garry Branch Royal Canadian Legion Poppy Trust Fund
Manitoba Centre for Nursing and Health Research for assistance with poster preparation



Background

- Estimates indicate that 20-50% of older people have undetected reduced vision, most of which are correctable or treatable (Smeeth et al, 2003).
- Rates of eye disease and visual impairment among nursing home residents is 3.3 times greater than any other segment of the population (Morer, 1994).
- Vision plays an important role in balance, mobility, falls and cognition in older persons.
- Falls are a major source of death and injury in the elderly (Harwood et al, 2005).
- Hip fractures are the most common fracture, the most devastating and the most costly to the health care system to treat (Kannus & Khan, 2001).
- In Manitoba, of the \$819 million per year spent on unintentional injuries, \$335 million is related to falls with \$164 million devoted to treating falls among the elderly (Papadimitropoulos et al., 1997)

Current Vision Care Guidelines

- Canada and Manitoba:
 - No specific guidelines
 - Vision has not been deemed a "medically necessary health service"
 - Lack of access and referral in the long term care setting

Objectives

- To identify prevalence of vision disorders.
- To test reliability and validity of the vision screening tool.
- To compare prevalence of falls, fractures, and various quality of life characteristics before and after treatment intervention.

Definitions

- Interventions: eye-specific vitamins, new eyeglass prescriptions, cataract surgery, eye drops, photodynamic therapy, avastin therapy, lighting improvements, and magnifiers
- Vision screening tool indicates:
 - Whether a person may benefit from a full eye exam from an optometrist, an ophthalmologist or general practitioner
 - Problems with vision that may be affecting day-to-day activities.
 - Detects vision loss caused from refractive errors, cataracts, macular degeneration, glaucoma and diabetic retinopathy
 - Vision Screening Tool: adapted from the Centre of Eye Research Australia/ WHO Vision Screening Tool

Study Design

- Convenience sample of residents 65 years and older in one long term care facility
- All nursing home residents who participated in a vision screening and intervention program at a long term care facility were approached
- Initial screening done by nurse
- Screening tool detects vision loss and provides the algorithm for appropriate referral to professional for diagnosis & treatment
- After screening by nurse, all residents were assessed by optometrist who was blinded to screening outcome
- Intervention was provided by the appropriate professional after optometrist assessment
- MDS data and other data from screening records were utilized to obtain data on prevalence of falls, fractures, changes in depression, social engagement, balance and quality of life



Sample Characteristics

- Sample Characteristics N=92
 - Female n=76 (82.6%)
 - Age Range = 63-102
 - Vision diagnosis at admission n= 18 (19.5%)
 - Wore glasses n=69 (75%)
 - No recorded data on last eye exam or if one had taken place n=70 (75%)
 - Fell in past 6 months n=19 (20%)

Findings

Reliability and Validity of Tool

- Screening by Nurse found 77 with vision impairment
- Optometrist identified 79 with vision impairment
 - Amount of agreement between nurse and optometrist: 97.5% agreement
- Nurse referred 52 for ophthalmology and 25 for optometry
- Optometrist identified 49 for ophthalmology and 28 for optometry
 - Strength of association between nurse and optometrist assessment: .799 (excellent)

Prevalence of vision impairment

- 74 did not have a vision impairment diagnosis at admission or in other words, only 18 had a diagnosis
- Upon screening, 72 were found with vision impairment
- Results of those who had an intervention (n=17)
- Significant improvement in level of depression, social engagement, balance and quality of life

- No occurrences of falls or fractures
- No occurrences of critical incidents

Results of those who received no intervention (n=29)

- Falls occurred in 18
- Critical incidents occurred in 8
- Types of fractures: hip 5, wrist 2, ribs 1
- Deaths: 3

Reasons for not receiving intervention:

- Mobility and transfer difficulties
- Ophthalmologist/Resident/Family stated intervention not needed or wanted:
 - "don't use vision much in a nursing home".
 - "won't make any difference".
- Deceased
- No reason given

2 predictors of falls post-intervention

- If had a previous fall (O.R.= 3.5, 1.06 – 11.2)
- If refused or did not receive intervention (O.R.=3.09, 1.13 – 8.44)



Conclusion

- The screening tool was easily used by the nurses
- The screening tool was highly reliable and valid in detecting vision loss
- The screening tool had high validity in referring to appropriate professional
- Those who had intervention had no falls/fractures and displayed improvements in at least one or more MDS and quality of life assessment categories
- Those who had no intervention were at greater risk for falls and fractures than those who obtained intervention
- Vision care is a major gap in the care of older adults in nursing homes

